## Commutative Algebra

Due date: Friday, 02/11/2007, 14h00

**Exercise 4:** Let R be a ring. Obviously  $R \hookrightarrow R[x_1, \dots, x_n] : a \mapsto a$  is a ring homomorphism and thus makes  $R[x_1, \dots, x_n]$  an R-algebra.

- a. Show that  $R[x_1,\ldots,x_n]$  satisfies the following universal property: if  $(R',\phi)$  is any R-algebra and  $\alpha_1,\ldots,\alpha_n\in R'$  are given, then there is a unique R-algebra homomorphism  $\alpha:R[x_1,\ldots,x_n]\to R'$  such that  $\alpha(x_i)=\alpha_i$  for all  $i=1,\ldots,n$ .
- b. Let  $I \subseteq R[x_1, \dots, x_n]$  and  $J \subseteq R[y_1, \dots, y_m]$ . Show that the following are equivalent:
  - (a)  $\phi:R[x_1,\ldots,x_n]/I\to R[y_1,\ldots,y_m]/J$  is an R-algebra homomorphism
  - (b) There are  $f_1, \ldots, f_n \in R[y_1, \ldots, y_m]$  such that  $g(f_1, \ldots, f_n) \in J$  for all  $g \in I$  and  $\phi(\overline{g}) = \overline{g(f_1, \ldots, f_n)}$  for all  $\overline{g} \in R[x_1, \ldots, x_n]/I$ .
  - (c) There is an R-algebra homomorphism  $\psi: R[x_1,\ldots,x_n] \to R[y_1,\ldots,y_m]$  such that  $\psi(I) \subseteq J$  and  $\varphi(\overline{g}) = \overline{\psi(g)}$ .

Note, a. means: we may uniquely define an R-algebra homomorphism on  $R[x_1, \ldots, x_n]$  by just specifying the images of the  $x_i$ !

**Exercise 5:** Let R be a ring and  $I, J_1, ..., J_n \subseteq R$ . Show that:

a. 
$$I:(\sum_{i=1}^n J_i) = \bigcap_{i=1}^n (I:J_i)$$
.

b. 
$$\left(\bigcap_{i=1}^n J_i\right): I = \bigcap_{i=1}^n (J_i: I)$$
.

c. 
$$\sqrt{J_1 \cap \ldots \cap J_n} = \sqrt{J_1} \cap \ldots \cap \sqrt{J_n}$$
.

$$d. \ \sqrt{J_1+\ldots+J_n} \supseteq \sqrt{J_1}+\ldots+\sqrt{J_n}.$$

**Exercise 6:** Let R be a ring and  $f = \sum_{n=0}^{\infty} a_n x^n \in R[[x]]$  a formal power series over R. Show:

- a. f is a unit if and only if  $a_0$  is a unit in R.
- b. What are the units in K[[x]] if K is a field?
- c. x is not a zero-divisor in R[[x]].
- d. If f is nilpotent, then  $a_n$  is nilpotent for all n. Is the converse true?