Fachbereich Mathematik Thomas Markwig Winter Semester 2007/08, Set 7 Henning Meyer

## **Commutative Algebra**

Due date: Friday, 07/12/2007, 14h00

**Exercise 24:** Let  $S \subseteq R$  be a multiplicatively closed subset, and consider the ring extension  $i: R \to S^{-1}R: r \mapsto \frac{r}{1}$ . Show that

 $\left\{ P \in \operatorname{\mathbf{Spec}}(R) \ \middle| \ S \cap P = \emptyset \right\} \longrightarrow \operatorname{\mathbf{Spec}}\left(S^{-1}R\right) : P \mapsto P^e = S^{-1}P$ 

is bijective with inverse

$$\operatorname{Spec}(S^{-1}R) \longrightarrow \left\{ P \in \operatorname{Spec}(R) \mid S \cap P = \emptyset \right\} : Q \mapsto Q^{c} = \mathfrak{i}^{-1}(Q).$$

In particular, for prime ideals  $P \in \text{Spec}(R)$  we have  $(P^e)^c = P$ .

## **Exercise 25:**

- a. Let K be a field,  $R = K[x, y, z]/\langle xz, yz \rangle$  and  $P = \langle x, y, z-1 \rangle \subseteq R$ . Show  $R_P \cong K[z]_{\langle z-1 \rangle}$ .
- b. Let R be a ring,  $f \in R$  a non-zero-divisor. Show  $R_f \cong R[x]/\langle fx 1 \rangle$ .

**Exercise 26:** Let R be a ring and  $\mathcal{N}(R)$  its nilradical. Show:

- a. If  $S \subseteq R$  multiplicatively closed, then  $\mathcal{N}(S^{-1}R) = S^{-1}\mathcal{N}(R)$ .
- b. A ring is called *reduced* if it has no nilpotent elements except 0. Show that "being reduced" is a local property, i.e. the following are equivalent:
  - (1) R is reduced.
  - (2)  $R_P$  is reduced for each  $P \in \text{Spec}(R)$ .
  - (3)  $R_m$  is reduced for each  $m \triangleleft \cdot R$ .
- c. Show that "being flat" is a local property, i.e. if M is an R-module, then the following are equivalent:
  - (1) M is a flat R-module.
  - (2)  $M_P$  is a flat  $R_P$  module for each  $P \in \text{Spec}(R)$ .
  - (3)  $M_{\mathfrak{m}}$  is a flat  $R_{\mathfrak{m}}$  module for each  $\mathfrak{m} \lhd \cdot R$ .

Hint for part c., use Exercise 21 and note that any  $R_P$ -module N is also an R-module and that  $N_P = N$ .

**Exercise 27:** Let  $I := \langle 2, 1 + \sqrt{-5} \rangle \triangleleft \mathbb{Z}[\sqrt{-5}]$ . Show that I as an R-module is projective, but not free.

Hint, note that  $2 \in I \cdot I$ . Use this to show that  $I \neq \langle x \rangle$  for any x, while for any prime P containing I we have  $I_P$  is generated by  $1 + \sqrt{-5}$ . For the last statement use Nakayama's Lemma in a sensible way!