

Commutative Algebra

Due date: Monday, 28/11/2011, 10h00

Exercise 21: Let $S \subseteq R$ be a multiplicatively closed subset, and consider the ring extension $i : R \rightarrow S^{-1}R : r \mapsto \frac{r}{1}$. Show that

$$\{P \in \text{Spec}(R) \mid S \cap P = \emptyset\} \longrightarrow \text{Spec}(S^{-1}R) : P \mapsto P^e = S^{-1}P$$

is bijective with inverse

$$\text{Spec}(S^{-1}R) \longrightarrow \{P \in \text{Spec}(R) \mid S \cap P = \emptyset\} : Q \mapsto Q^c = i^{-1}(Q).$$

In particular, for prime ideals $P \in \text{Spec}(R)$ we have $(P^e)^c = P$.

Exercise 22:

- Let K be a field, $R = K[x, y, z]/\langle xz, yz \rangle$ and $P = \langle x, y, z - 1 \rangle \trianglelefteq R$. Show $R_P \cong K[z]_{\langle z-1 \rangle}$.
- Let R be a ring, $f \in R$ a non-zero-divisor. Show $R_f \cong R[x]/\langle fx - 1 \rangle$.

Exercise 23: Let R be a ring and $\mathcal{N}(R)$ its nilradical. Show:

- If $S \subseteq R$ multiplicatively closed, then $\mathcal{N}(S^{-1}R) = S^{-1}\mathcal{N}(R)$.
- A ring is called *reduced* if it has no nilpotent elements except 0. Show that “being reduced” is a local property, i.e. the following are equivalent:
 - R is reduced.
 - R_P is reduced for each $P \in \text{Spec}(R)$.
 - R_m is reduced for each $m \triangleleft \cdot R$.
- Show that “being flat” is a local property, i.e. if M is an R -module, then the following are equivalent:
 - M is a flat R -module.
 - M_P is a flat R_P module for each $P \in \text{Spec}(R)$.
 - M_m is a flat R_m module for each $m \triangleleft \cdot R$.

Hint for part c., use Exercise 21 and note that any R_P -module N is also an R -module and that $N_P = N$.

Exercise 24: Let $I := \langle 2, 1 + \sqrt{-5} \rangle \triangleleft \mathbb{Z}[\sqrt{-5}]$. Show that I as an R -module is projective, but not free.

Hint, note that $2 \in I \cdot I$. Use this to show that $I \neq \langle x \rangle$ for any x , while for any prime P containing I we have I_P is generated by $1 + \sqrt{-5}$. For the last statement use Nakayama’s Lemma in a sensible way!

In-Class Exercise 13: Let $R = K[x, y]$ and $P = K[x, y, z]/\langle xz - x, yz - y - z + 1 \rangle$. Is P a flat R -module?

In-Class Exercise 14: Let $m = \langle x, y \rangle \triangleleft K[x, y]$. Localize the ring R and the module P in In-Class Exercise 13 at m . Is the resulting module P_m a flat R_m -module?