

## Commutative Algebra

Due date: Monday, 09/02/2015, 10h00

**Exercise 48:** Let  $K$  be any field, and  $\underline{\alpha} = (\alpha_1, \dots, \alpha_n) \in \mathbb{R}^n$  be an independent set of real numbers. Show:

- $\varphi_{\underline{\alpha}} : K(x_1, \dots, x_n) \rightarrow K\{\{t\}\} : \frac{f}{g} \mapsto \frac{f(t^{\alpha_1}, \dots, t^{\alpha_n})}{g(t^{\alpha_1}, \dots, t^{\alpha_n})}$  is a  $K$ -algebra monomorphism.
- $v : K(x_1, \dots, x_n)^* \mapsto \mathbb{R} : h \mapsto (\text{ord} \circ \varphi_{\underline{\alpha}})(h)$  is a valuation of  $K(x_1, \dots, x_n)$ .
- $1 = \dim(R_v) < \text{trdeg}_K(K(x_1, \dots, x_n)) - \text{trdeg}_K(R_v/\mathfrak{m}_{R_v}) = n$ , for  $n \geq 2$ .

Note,  $\text{ord} : K\{\{t\}\}^* \rightarrow \mathbb{R}$  is the valuation of  $K\{\{t\}\}$  from Exercise 50.

**Exercise 49:** Let  $R$  be a Dedekind domain and  $0 \notin S \subset R$  multiplicatively closed. Show that either  $S^{-1}R = \text{Quot}(R)$  or  $S^{-1}R$  is a Dedekind domain.

**Exercise 50: [Lemma of Gauß]\***

Let  $R$  be a Dedekind domain. For a polynomial  $f = \sum_{i=0}^n a_i x^i \in R[x]$  we call  $c(f) = \langle a_0, \dots, a_n \rangle_R$  the *content* of  $f$ . Show that  $c(f) \cdot c(g) = c(f \cdot g)$  for  $f, g \in R[x]$ .

Hint, reduce to the case that  $R$  is local (i.e. a DVR), and use Nakayama's Lemma in a suitable way.

**Exercise 51: [Chinese Remainder Theorem]**

Let  $R$  be a Dedekind domain and  $I_1, \dots, I_n \leq R$ .

- Show that the following sequence is exact

$$R \xrightarrow{\varphi} \bigoplus_{i=1}^n R/I_i \xrightarrow{\psi} \bigoplus_{i < j} R/(I_i + I_j),$$

where  $\varphi(x) = (x + I_1, \dots, x + I_n)$  and  $\psi(x_1 + I_1, \dots, x_n + I_n) = (x_i - x_j + I_i + I_j)_{i < j}$ .

- Given  $x_1, \dots, x_n \in R$ . Show there is an  $x \in R$  such that  $x \equiv x_i \pmod{I_i}$  for  $i = 1, \dots, n$  if and only if  $x_i \equiv x_j \pmod{I_i + I_j}$  for  $i \neq j$ .

Hint for part a., localize with respect to maximal ideals! – Note, part b. generalizes 1.12.

**In-Class Exercise 27:** Describe  $\text{Div}(R)$  and  $\text{Pic}(R)$  for  $R = K[x, y]/\langle y - x^2 \rangle$ ?

---

\*What is the connection to the *Lemma of Gauß* in 1.38, stating “ $R$  factorial implies  $R[x]$  factorial”? If we replace the assumption “ $R$  Dedekind domain” by “ $R$  UFD” the above result holds true as well. Call a polynomial *primitive* if  $c(f) = R$  (or equivalently if  $R^*$  is the gcd of the coefficients of  $f$ ), then we deduce from the above result that a primitive polynomial in  $R[x]$  can only factorize in a product of primitive polynomials, which are then necessarily of smaller degree. By induction on the degree we see that each primitive polynomial is a product of irreducible primitive polynomials. Thus, every polynomial is a product of irreducible ones, since splitting off a greatest common divisor  $g$  of its coefficients gives a primitive one and  $g$  factorises since  $R$  is factorial. – It then only remains to show that each irreducible polynomial in  $R[x]$  is prime. – In the literature it is more common to call the statement “ $R$  UFD implies  $c(f \cdot g) = c(f) \cdot c(g)$ ” the *Lemma of Gauß*.