

Computer Algebra

Due date: Tuesday, 20/11/2007, 10h00

Exercise 9: Let $I \trianglelefteq K[x]$ be a homogeneous ideal in the polynomial ring. Show that the degree reverse lexicographical ordering $>_{dp}$ satisfies the properties:

$$L(I + x_n^d) = L(I) + \langle x_n^d \rangle \quad \text{and} \quad L(I : x_n^d) = L(I) : x_n^d \quad \text{for any } d \geq 1.$$

Note, since I is homogeneous we only have to consider homogeneous polynomials – e. g. $L(I) = \langle \text{lm}(f) \mid f \in I, f \text{ homogeneous} \rangle$, etc.

Exercise 10: Apply IDBUCHBERGER to the following data (without using Singular):

$$g = x^4 + y^4 + z^4 + xyz, G = \left(\frac{\partial g}{\partial x}, \frac{\partial g}{\partial y}, \frac{\partial g}{\partial z} \right), >_{dp}.$$

Exercise 11: Write a SINGULAR procedure `IDBuchberger` which takes as input a list consisting of a polynomial g and a list of polynomials f_1, \dots, f_k and returns an indeterminate division with remainder of g with respect to (f_1, \dots, f_k) in the form of a list consisting of the remainder r and a list of the scalars q_1, \dots, q_k .

Note, the ordering of the base ring must be global! For a list of polynomials it is best to use the type `ideal`.

Exercise 12: Write a SINGULAR procedure `RedIDBuchberger` which takes as input a list consisting of a polynomial g and a list of polynomials f_1, \dots, f_k and returns a reduced indeterminate division with remainder of g with respect to (f_1, \dots, f_k) in the form of a list consisting of the remainder r and a list of the scalars q_1, \dots, q_k . I suggest that you do your implementation in such a way that you actually get a determinate division with remainder.

Note, the ordering of the base ring must be global!