

## Computer Algebra

Due date: Tuesday, 18/12/2007, 10h00

**Exercise 25:** Let  $>$  be a global monomial ordering on  $\text{Mon}(\underline{x})$ , and let  $M \in \text{Mat}(n \times n, K[\underline{x}])$ . By  $f_1, \dots, f_n \in K[\underline{x}]^{2n}$  we denote the rows of the matrix  $(M, \mathbb{1}_n)$ , and  $G = (g_1, \dots, g_k)$  shall be *the* reduced standard basis of  $\langle f_1, \dots, f_k \rangle_{K[\underline{x}]} \leq K[\underline{x}]^{2n}$  w. r. t. the ordering  $(c, >)$  with  $\text{lm}(g_1) > \dots > \text{lm}(g_k)$ . Show:

- $M$  is invertible if and only if  $k = n$  and  $\text{lm}(g_i) = e_i$  for  $i = 1, \dots, n$ .
- If  $M$  is invertible, then the rows of  $(\mathbb{1}_n, M^{-1})$  are just  $g_1, \dots, g_n$ .

**Exercise 26:** Let  $R = \mathbb{Q}[x, y, z]/\langle x^2 + y^2 + z^2 \rangle$ ,  $M = R^3/\langle (x, xy, xz)^t \rangle$  and  $N = R^2/\langle (1, y)^t \rangle$ . Moreover, let  $\varphi : M \rightarrow N$  be given by the matrix

$$A = \begin{pmatrix} x^2 + 1 & y & z \\ yz & 1 & -y \end{pmatrix}.$$

- Compute  $\text{Ker}(\varphi)$ .
- Test if  $(x^2, y^2)^t \in \text{Im}(\varphi)$ .
- Compute  $\text{Im}(\varphi) \cap \{f \in N \mid f \equiv (h, 0) \text{ mod } \langle (x, 1)^t \rangle \text{ for some } h \in R\}$ .
- Compute  $\text{ann}_R(\text{Im}(\varphi))$ .

Note, you may use Singular for your computations!

**Exercise 27:** Write a SINGULAR procedure `intersection` which takes as input two lists consisting of polynomials  $f_1, \dots, f_k$  respectively  $g_1, \dots, g_l$  and returns a standard basis of the ideal  $\langle f_1, \dots, f_k \rangle \cap \langle g_1, \dots, g_l \rangle$ .

**Exercise 28:** Write a SINGULAR procedure `idealquotient` which takes as input two lists consisting of polynomials  $f_1, \dots, f_k$  respectively  $g_1, \dots, g_l$  and returns a standard basis of the ideal  $\langle f_1, \dots, f_k \rangle : \langle g_1, \dots, g_l \rangle$ .