

## Introduction to Commutative Algebra and Algebraic Geometry

### Exercise Sheet 3

---

#### Exercise 1.

Let  $X_1 = V(x^3 - 3x + 2 - y^2)$ ,  $X_2 = V(x - 1)$  be two varieties in  $\mathbb{C}^2$ . Make a sketch of the real part of  $X_1, X_2 \subset \mathbb{R}^2$  in the same coordinate system and compute  $\sqrt{I(X_1) + I(X_2)} \subset \mathbb{C}[X, Y]$ .

#### Exercise 2.

Let  $K$  be a field. Consider the variety  $V = V(Y^2 - X^3)$  of the cuspidal cubic  $Y^2 - X^3 \in K[X, Y]$ . Let  $K[V]$  be the coordinate ring of  $V$ . Prove

$$K[V] \cong K[T^2, T^3].$$

#### Exercise 3.

Let  $K$  be an algebraically closed field. Find all polynomials  $f \in K[X_1, \dots, X_n]$  for which  $V(X_{n+1}^2 - f) \subset K^{n+1}$  is irreducible. Prove your claim.

#### Exercise 4.

Let  $K$  be a field and let  $0 \rightarrow V_0 \rightarrow \dots \rightarrow V_n \rightarrow 0$  be an exact sequence of finite dimensional vector spaces over  $K$ . Prove:

$$\sum_{i=0}^n (-1)^i \dim(V_i) = 0.$$