

## MATHEMATICAL STATISTICAL PHYSICS: ASSIGNMENT 10

**Problem 42:** *Moments of a random wave function* (hand in, 50 points)

Let  $\mathbb{S}$  be the unit sphere in the Hilbert space  $\mathbb{C}^d$ ,  $u$  the uniform probability distribution on  $\mathbb{S}$ , and  $\Psi = (\Psi_1, \dots, \Psi_d) \sim u$ . Compute all moments of  $\Psi$  of up to fourth order. That is, show for all  $k, \ell, m, n \in \{1, \dots, d\}$  that

- (a)  $\mathbb{E}\Psi_k = 0$  (*Hint:* symmetry)
- (b)  $\mathbb{E}\Psi_k^* \Psi_\ell = 0 = \mathbb{E}\Psi_k \Psi_\ell$  for  $k \neq \ell$
- (c)  $\mathbb{E}|\Psi_k|^2 = 1/d$
- (d)  $\mathbb{E}\Psi_k^2 = 0$
- (e)  $\mathbb{E}\Psi_k \Psi_\ell \Psi_m = 0$ , and likewise if any of the factors is conjugated
- (f)  $\mathbb{E}\Psi_k \Psi_\ell \Psi_m \Psi_n = 0$  if an index occurs only once, and likewise for conjugated factors
- (g)  $\mathbb{E}\Psi_k^4 = 0 = \mathbb{E}\Psi_k^{*4} = \mathbb{E}\Psi_k^* \Psi_k^3 = \mathbb{E}\Psi_k^{*3} \Psi_k$
- (h)  $\mathbb{E}|\Psi_k|^4 = \frac{2}{d(d+1)}$  (the main problem!)

(*Instructions:* Regard  $\mathbb{C}^d$  as  $\mathbb{R}^{2d}$ ,  $\Psi = (x_1, \dots, x_{2d}) = \mathbf{x}$ ,  $I_1 = \int_{\mathbb{S}} u(d\mathbf{x}) x_1^4$ ,

$I_2 = \int_{\mathbb{S}} u(d\mathbf{x}) x_1^2 x_2^2$ . Integrating in spherical coordinates,<sup>1</sup>

$$\int_{\mathbb{R}^{2d}} d\mathbf{x} x_1^2 x_2^2 \exp(-|\mathbf{x}|^2) = \int_0^\infty dr r^{2d-1} r^4 \exp(-r^2) I_2 \text{area}(\mathbb{S}). \quad (1)$$

Now the substitution  $s = r^2$  helps. Use without proof that  $\mathbb{E}[(X - \mu)^4] = 3\sigma^4$  for  $X \sim \mathcal{N}(\mu, \sigma^2)$ .

- (i)  $\mathbb{E}|\Psi_k|^2 |\Psi_\ell|^2 = \frac{1}{d(d+1)}$  for  $k \neq \ell$  (*Hint:*  $\mathbb{E}[(\sum_k |\Psi_k|^2)^2] = 1$  (why?).)
- (j)  $\mathbb{E}\Psi_k^2 \Psi_\ell^2 = 0 = \mathbb{E}|\Psi_k|^2 \Psi_\ell^2 = \mathbb{E}\Psi_k^{*2} \Psi_\ell^2$  for  $k \neq \ell$ .

**Problem 43:** *Variance and covariance of a random wave function* (hand in, 25 points)

(a) For  $\Psi$  as in Problem 42, conclude from the results of Problem 42 that

$$\text{Var}(|\Psi_1|^2) = \frac{1}{d^2} \frac{d-1}{d+1}, \quad \text{Cov}(|\Psi_1|^2, |\Psi_2|^2) = -\frac{1}{d^2(d+1)}.$$

(b) As we know, for large  $d$ ,  $\Psi_1$  is approximately  $\mathcal{N}^2(\mathbf{0}, I/2d)$  distributed. For comparison, let  $\mathbf{G} = (G_1, \dots, G_d) = (X_1, \dots, X_{2d})$  be a Gaussian random vector in  $\mathbb{C}^d = \mathbb{R}^{2d}$ , i.e., so that the  $X_i$  (the real and imaginary parts of the  $G_k$ ) are i.i.d. with  $X_i \sim \mathcal{N}(0, 1/2d)$ . Determine  $\text{Var}(|G_1|^2)$  and  $\text{Cov}(|G_1|^2, |G_2|^2)$ .

<sup>1</sup>This trick was discovered by N. Ullah, *Nuclear Physics* **58**: 65–71 (1964).

**Problem 44:** *Quantum particle in a box in 1d* (don't hand in)

(a) On the interval  $[0, L]$ , consider the Hamiltonian operator  $H\psi(x) = -\psi''(x)/2m$  with Dirichlet boundary conditions  $\psi(0) = 0$ ,  $\psi(L) = 0$ . Verify that the normalized eigenfunctions read

$$\varphi_n(q) = \left(\frac{2}{L}\right)^{1/2} \sin(n\frac{\pi}{L}q) \quad (2)$$

with  $n \in \mathbb{N}$  and eigenvalues

$$E_n = \frac{\pi^2}{2mL^2} n^2. \quad (3)$$

(b) It is known from Fourier series that the functions  $1, \sin nx, \cos nx$  ( $n \in \mathbb{N}$ ), after normalization, form an orthonormal basis of  $L^2([-\pi, \pi])$ . How can we conclude that the functions (2) form an orthonormal basis of  $L^2([0, L])$ ?

**Problem 45: Projection to fermionic wave functions** (hand in, 25 points)

We write  $(-1)^\sigma$  for the sign of a permutation  $\sigma \in S_N$ . We want to show that  $P_-$  defined by

$$P_- \psi(\mathbf{q}_1, \dots, \mathbf{q}_N) = \frac{1}{N!} \sum_{\sigma \in S_N} (-1)^\sigma \psi(\mathbf{q}_{\sigma(1)}, \dots, \mathbf{q}_{\sigma(N)}) \quad (4)$$

is the orthogonal projection to the subspaces of anti-symmetric functions in  $\mathcal{H} = L^2(\mathbb{R}^{3N})$ . Proceed as follows:

- (a)  $P_- \psi$  is an anti-symmetric function.
- (b) If  $\psi$  is already anti-symmetric, then  $P_- \psi = \psi$ .
- (c)  $P_-^2 = P_-$
- (d)  $P_- : \mathcal{H} \rightarrow \mathcal{H}$  is self-adjoint.

From (c) and (d) it follows that  $P_-$  is an orthogonal projection, and from (a) and (b) that its range is the space of anti-symmetric functions in  $\mathcal{H}$ .

**Hand in:** By 8:15am on Tuesday, July 5, 2022 via [urm.math.uni-tuebingen.de](http://urm.math.uni-tuebingen.de)