

## FOUNDATIONS OF QUANTUM MECHANICS: ASSIGNMENT 1

**Exercise 1: Essay question.** What got you interested in the foundations of quantum mechanics?

**Exercise 2: Plane waves.**

Show that for every constant vector  $\mathbf{k} \in \mathbb{R}^3$  (the *wave vector*), there is a unique constant  $\omega \in \mathbb{R}$  so that

$$\psi(t, \mathbf{x}) = e^{i\mathbf{k} \cdot \mathbf{x}} e^{-i\omega t} \quad (1)$$

satisfies the free (i.e.,  $V = 0$ ) Schrödinger equation (2.1) for  $N = 1$ . Specify  $\omega$  in terms of  $\mathbf{k}$ . Remark: Since for every  $t$ ,  $\|\psi_t\| = \infty$ , this function (called a plane wave) is not square-integrable (not normalizable) and thus not physically possible as a wave function; but it is a useful toy example.

**Exercise 3: Time reversal invariance.**

Show that if  $\psi(t, q)$  is a solution of the Schrödinger equation (2.1) with real-valued potential  $V$ , then so is  $\psi^*(-t, q)$ .

**Exercise 4: Gaussian wave packet.**

(a) Show that the function

$$\psi(\mathbf{x}, t) = (2\pi\lambda_t^2\sigma^2)^{-3/4} e^{i\mathbf{k} \cdot (\mathbf{x} - \hbar\mathbf{k}t/2m)} e^{-\frac{(\mathbf{x} - \hbar\mathbf{k}t/m)^2}{4\lambda_t\sigma^2}} \quad (2)$$

with

$$\lambda_t = 1 + \frac{i\hbar t}{2m\sigma^2} \quad (3)$$

and arbitrary constants  $\mathbf{k} \in \mathbb{R}^3$ ,  $\sigma > 0$ , is a solution of the free Schrödinger equation of a single particle in 3 dimensions,

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi. \quad (4)$$

The main difficulty here is to organize the calculation so as to make it manageable.

(b) Show that the probability density  $\rho_t(\mathbf{x})$  is, at every  $t$ , Gaussian and specify its mean and standard deviation.

**Hand in:** Tuesday October 22, 2019, in class

---

**Reading assignment** due Friday October 25, 2019: R. Feynman, *Feynman Lectures on Physics* vol. 3, chapter 1.