Universität Tübingen, Fachbereich Mathematik Dr. Stefan Keppeler & Saradha Senthil-Velu

## Groups and Representations

Homework Assignment 6 (due on 30 November 2022)

## Problem 21

We consider a rotationally invariant Hamiltonian. Let E be an eigenvalue of H with eigenspace  $V_E$  spanned by the spherical harmonics  $Y_{1m}(\varphi, \vartheta) = \cos \vartheta e^{im\varphi}$  with a fixed radial part R, i.e.  $V_E = \operatorname{span}(\{R(r)Y_{1m}(\varphi, \vartheta) : m = -1, 0, 1\})^2$ 

 $V_E$  carries a three-dimensional irreducible representation of O(3), defined by  $(\Gamma(U)\psi)(x) = \psi(U^{-1}x)$ . O(3) contains the subgroup  $D_3 = \{e, C, \overline{C}, \sigma_1, \sigma_2, \sigma_3\} \cong S_3$ , where C and  $\overline{C}$  denote rotations about the z-axis (cf. Section 2.4.1).

Study the effect of perturbations that are only invariant under  $D_3$  or  $\mathbb{Z}_3 \cong \{e, C, \overline{C}\}$ . Determine the relevant irreducible representations with their multiplicities and sketch the possible splitting of energy levels.

## Problem 22

We consider once more the  $CO_2$  molecule of Problems 10 & 18. In Problem 10 we found a six-dimensional representation of  $V_4$ . Decompose the six-dimensional carrier space into irreducible invariant subspaces by applying the generalised projection operators.

<sup>2</sup>We use spherical coordinates

 $\begin{pmatrix} x\\ y\\ z \end{pmatrix} = \begin{pmatrix} r\sin\vartheta\cos\varphi\\ r\sin\vartheta\sin\varphi\\ r\cos\vartheta \end{pmatrix}.$ 

## Problem 23

For  $\sigma \in S_n$  and j = 1, ..., n let  $k_j(\sigma)$  be the number of (disjoint) cycles of length j in  $\sigma$ , e.g.  $k_1(e) = n$  and  $k_j(e) = 0 \forall j > 1$ . Show:

a) The conjugacy class of  $\sigma$  is determined by its cycle structure, i.e.

$$[\sigma] := \{\tau \sigma \tau^{-1} : \tau \in S_n\} = \{\tau \in S_n : k_j(\tau) = k_j(\sigma), j = 1, \dots, n\}$$

It's almost trivial using the birdtrack notation (see Section 1.4)!

HINT: In order to make the cycle structure visible consider the birdtrack diagram of  $\sigma$  and connect the first line on the left to the first line on the right etc.; e.g. for  $(12), (132) \in S_3$  consider



b) The number of elements of a class is given by

$$|[\sigma]| = rac{n!}{\prod\limits_{j \le n} k_j! j^{k_j}}$$
 .

c) Fun exercise (optional): Watch the video An Impossible Bet by minutephysics,

https://youtu.be/eivGlBKlK6M,

and come up with a good strategy. Don't watch the solution! Think about cycles instead.