

## Groups and Representations

Homework Assignment 9 (due on 21 December 2022)

---

### Problem 32

Verify that the (suitably normalised) Young operators for the standard Young tableaux for  $S_3$  (see Section 5.3) add up to the identity.

### Problem 33

For a fixed partition  $\lambda$  of  $n \in \mathbb{N}$  we define an ordering of standard tableaux as follows. Consider the sequence  $(r_\lambda^p)_j, j = 1, \dots, n$ , of numbers in the boxes of  $\Theta_\lambda^p$  starting with the first row read from left to right, then the second row from left to right etc. We say that  $\Theta_\lambda^p > \Theta_\lambda^q$  if the first non-vanishing term in the sequence  $(r_\lambda^p)_j - (r_\lambda^q)_j, j = 1, \dots, n$ , is positive. Then, e.g., the standard tableaux for  $\lambda = (3, 2)$  are ordered as

$$\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 4 & 5 & \\ \hline \end{array} < \begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 3 & 5 & \\ \hline \end{array} < \begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & \\ \hline \end{array} < \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 5 & \\ \hline \end{array} < \begin{array}{|c|c|c|} \hline 1 & 3 & 5 \\ \hline 2 & 4 & \\ \hline \end{array} .$$

- a) Prove that  $e_\lambda^p e_\lambda^q = 0$  if  $\Theta_\lambda^p > \Theta_\lambda^q$ .
- b) Show that (a) implies that the left ideals generated by the standard tableaux for a fixed partition are linearly independent.

### Problem 34

- a) Determine the character table of  $S_4$  using the methods of Section 5.5.
- b) Consider the following product representations of  $S_4$ , determine which irreps they contain and how many times. Also write their decomposition into irreps in terms of Young diagrams.

$$\begin{array}{ll} \text{(i)} & \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \\ \hline \end{array} \otimes \begin{array}{|c|c|c|} \hline \\ \hline \\ \hline \end{array} & \text{(ii)} & \begin{array}{|c|c|c|} \hline \\ \hline \\ \hline \end{array} \otimes \begin{array}{|c|} \hline \\ \hline \\ \hline \\ \hline \end{array} \\ \text{(iii)} & \begin{array}{|c|c|c|} \hline \\ \hline \\ \hline \end{array} \otimes \begin{array}{|c|c|} \hline \\ \hline \\ \hline \end{array} & \text{(iv)} & \begin{array}{|c|c|} \hline \\ \hline \\ \hline \end{array} \otimes \begin{array}{|c|c|} \hline \\ \hline \\ \hline \end{array} \end{array}$$

### Problem 35

In the quark model baryons are made out of three quarks. The latter are characterised i.a. by the quantum numbers<sup>3</sup>  $I$  (isospin) and  $Y$  (hyper charge).

We have  $(I, Y) = (\frac{1}{2}, \frac{1}{3})$  for the up-quark,  $|u\rangle$ ,  $(I, Y) = (-\frac{1}{2}, \frac{1}{3})$  for the down-quark,  $|d\rangle$ , and  $(I, Y) = (0, -\frac{2}{3})$  for the strange-quark,  $|s\rangle$ . For products like  $|udd\rangle = |u\rangle \otimes |d\rangle \otimes |d\rangle$  the values of  $I$  and  $Y$  are given by the sums of the values for the individual quarks.

For combinations of 3 quarks (each up, down or strange) we thus have a 27-dimensional space  $V$ , which carries a representation of  $S_3$  (by permutation of the factors).

- Which irreps are contained in this representation and what are their multiplicities?
- Let  $U \subset V$  be an irreducible invariant subspace. What can we say about the values of  $I$  and  $Y$  on  $U$ ?
- In a  $(I, Y)$ -diagram mark all points corresponding to vectors transforming in the irrep defined by  $\square\square\square$ .
- Repeat part (c) for the irrep with Young diagram  $\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$ . You find some potentially useful Octave/MATLAB-Code on the course webpage.

---

<sup>3</sup>Think of “quantum number” as “eigenvalue of some linear operator”.