

## FOUNDATIONS OF QUANTUM MECHANICS

Written homework due Wednesday November 22, 2017

**Exercise 9: Essay question.** Explain what reductionism means.

### Exercise 10: Fourier transform

(a) Find the Fourier transform  $\widehat{\psi}$  of the function

$$\psi(x) = \begin{cases} 0 & x < -1 \\ 2^{-1/2} & -1 \leq x \leq 1 \\ 0 & x > 1, \end{cases} \quad (1)$$

where  $x$  is a 1-dimensional variable.

(b) (optional) Plot  $\widehat{\psi}$  using a computer.

(c) (optional) Using suitable software (such as Mathematica, Maple, or Matlab), make the computer find the formula for  $\widehat{\psi}$ . (That is, you need to find the command for symbolically computing Fourier transforms and the one for defining a function like (1), and run them.)

### Exercise 11: Pauli matrices

The three *Pauli matrices* are

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2)$$

(a) For each of  $\sigma_1$  and  $\sigma_2$ , find an ONB of eigenvectors in  $\mathbb{C}^2$ .

(b) Show that for every unit vector  $\mathbf{n} \in \mathbb{R}^3$ , the Pauli matrix in direction  $\mathbf{n}$ ,  $\sigma_{\mathbf{n}} := \mathbf{n} \cdot \boldsymbol{\sigma} = n_1\sigma_1 + n_2\sigma_2 + n_3\sigma_3$ , has eigenvalues  $\pm 1$ . (Hint: compute det and trace.)

(c) Show that every self-adjoint complex  $2 \times 2$  matrix  $A$  is of the form  $A = cI + \mathbf{u} \cdot \boldsymbol{\sigma}$  with  $c \in \mathbb{R}$  and  $\mathbf{u} \in \mathbb{R}^3$ .

### Exercise 12: Potential step

Consider a potential step  $V(x) = V_0 1_{0 \leq x}$  in 1d with  $V_0 > 0$ . We want to compute the reflection and transmission probabilities as a function of the incoming momentum  $\hbar k_0$ , assuming that  $E := \hbar^2 k_0^2 / 2m > V_0$ . A recipe for that is to consider a plane wave  $e^{ik_0 x}$  coming from  $x = -\infty$  and see how much gets reflected and transmitted by constructing an eigenfunction  $\psi$  of  $H$ ,  $H\psi = E\psi$ , from the ansatz

$$\psi(x) = \begin{cases} Ae^{ik_0 x} + Be^{-ik_0 x} & \text{for } x < 0 \\ Ce^{i\kappa_0 x} & \text{for } x > 0 \end{cases} \quad (3)$$

with complex coefficients  $A, B, C$ , representing the incoming wave  $e^{ik_0 x}$ , the reflected wave  $e^{-ik_0 x}$ , and the transmitted wave  $e^{i\kappa_0 x}$ . Regarding  $V$  as a limit of continuous functions leads to the further requirement that  $\psi$  be continuous and continuously differentiable at 0.

(a) Determine  $\kappa_0$  from  $H\psi = E\psi$ .

(b) Determine  $B$  and  $C$  from  $A$ .

(c) The recipe says that the three waves are associated with probability currents  $j_{\text{in}} = \hbar k_0 |A|^2/m$ ,  $j_R = \hbar k_0 |B|^2/m$ , and  $j_T = \hbar \kappa_0 |C|^2/m$ ; and that the reflection probability is  $P_R = j_R/j_{\text{in}}$ , while the transmission probability is  $P_T = j_T/j_{\text{in}}$ . Compute  $P_R$  and  $P_T$ .

(d) To justify the recipe, consider a “plane wave packet”  $\psi(x) = \phi(x) e^{ik_0 x}$  with a (non-Gaussian) envelope profile  $\phi(x)$  that is nearly constant over a region of length  $L$  much larger than the wave length  $2\pi/k_0$  and then drops to 0 over a length much smaller than  $L$  but still much larger than the wave length. Let us take for granted that under the free Schrödinger evolution the envelope function  $\phi$  will approximately maintain its shape (in particular, its length) for a long time and simply move at speed  $\hbar k_0/m$ . Suppose the packet hits the step at  $t = 0$ ; forget about what happens at the edges of the packet and focus on the bulk. A reflected plane wave packet and a transmitted one are generated; at time  $\tau > 0$ , the incoming plane wave packet is used up, and the two outgoing packets end. During  $0 < t < \tau$ , the region around 0 is well approximated by (3); after  $\tau$ , the outgoing packets keep moving away from the origin. Determine  $\tau$  and the lengths  $L_R$  and  $L_T$  of the reflected and transmitted packets  $\psi_R$  and  $\psi_T$ .

(e) Determine  $P_R = \|\psi_R\|^2$  and  $P_T = \|\psi_T\|^2$  and verify that they agree with part (c).

(f) Draw a space-time diagram of the Bohmian trajectories in the bulk (that is, ignoring any edge effects).

---

**Reading assignment** due Wednesday November 29, 2017: T. Maudlin, Three Measurement Problems. *Topoi* **14(1)**: 7–15 (1995). Read pages 7–12 and the first two paragraphs on page 13.