

## Commutative Algebra

**In-Class Exercise 1:** We call an ideal  $I$  in the polynomial ring  $K[\underline{x}] = K[x_1, \dots, x_n]$  a *monomial ideal* if  $I$  is generated by (possibly infinitely many) monomials.

Given two monomials  $\underline{x}^\alpha$  and  $\underline{x}^\beta$  we say that  $\underline{x}^\alpha$  *divides*  $\underline{x}^\beta$  if there is a monomial  $\underline{x}^\gamma$  such that  $\underline{x}^\alpha \cdot \underline{x}^\gamma = \underline{x}^\beta$ , i.e.  $\alpha_i \leq \beta_i$  for all  $i = 1, \dots, n$ .

And we define the *least common multiple* of  $\underline{x}^\alpha$  and  $\underline{x}^\beta$  in the obvious way as

$$\text{lcm}(\underline{x}^\alpha, \underline{x}^\beta) = x_1^{\max\{\alpha_1, \beta_1\}} \dots x_n^{\max\{\alpha_n, \beta_n\}},$$

i.e. it is the monomial of lowest degree which is divisible by both monomials.

a. Show that for an ideal  $I$  the following are equivalent:

- (1)  $I$  is a monomial ideal.
- (2) For any  $f \in I$  also all monomials occurring in  $f$  belong to  $I$ .
- (3) There is a generating set  $B$  of  $I$  such that for any  $f \in B$  all monomials of  $f$  belong to  $I$ .

b. If  $I = \langle \underline{x}^\alpha \mid \alpha \in \Lambda \rangle$  and  $\underline{x}^\beta \in I$  then there is an  $\alpha \in \Lambda$  such that  $\underline{x}^\alpha$  divides  $\underline{x}^\beta$ .

c. Let  $I = \langle \underline{x}^\alpha \mid \alpha \in \Lambda \rangle$  and  $J = \langle \underline{x}^\beta \mid \beta \in \Lambda' \rangle$  be two monomial ideals in  $K[\underline{x}]$ . Show that

$$I \cap J = \langle \text{lcm}(\underline{x}^\alpha, \underline{x}^\beta) \mid \alpha \in \Lambda, \beta \in \Lambda' \rangle$$

and

$$I : \langle \underline{x}^\gamma \rangle = \left\langle \frac{\text{lcm}(\underline{x}^\alpha, \underline{x}^\gamma)}{\underline{x}^\gamma} \mid \alpha \in \Lambda \right\rangle.$$

Hint for part c., show first that the two ideals are monomial ideals.

**In-Class Exercise 2:** We will now introduce some basic commands for SINGULAR. In SINGULAR we have can work with two types of rings that we have introduced so far in the lecture, polynomial rings  $K[x_1, \dots, x_n]$  and power series rings  $K[[x_1, \dots, x_n]]$ . The polynomial ring  $\mathbb{Q}[x, y, z]$  is defined in SINGULAR as:

```
ring r=0, (x,y,z), dp;
```

Here, 0 stands for the characteristic of  $\mathbb{Q}$  and `dp` says that we are working with a **polynomial ring**.

The power series ring  $\mathbb{Z}/5\mathbb{Z}[[x_1, \dots, x_4]]$  is defined in SINGULAR as:

```
ring r=5,(x(1..4)),ds;
```

Here, 5 stands for the characteristic of  $\mathbb{Z}/5\mathbb{Z}$  and `dp` says that we are working with a power series ring — actually this is not quite true, but morally it is, and we need the notion of *localisation* to be more precise.

Once we have fixed a ring we can define polynomials and ideals and perform operations with them:

```
LIB "all.lib";          // load libraries needed e.g. for the radical
ring r=0,(x,y,z),dp;
poly f=x^3*y+5*z^2;
poly g=3x2y-xz2;      // this is short hand for 3*x^2*y-x*z^2
ideal I=f,g,x2y;
ideal J=x+y;
I*J;                  // the product of I and J
intersect(I,J);       // intersect the two ideals
quotient(I,J);        // compute the ideal quotient
radical(I);           // compute the radical of I
I=std(I);             // replace the generators of I by better ones
reduce(f,I);          // test if f belongs to I
reduce(J,I);          // test if J is contained in I
```

Consider the ideal  $I = \langle x^2y^5, x^6, y^2 \rangle$  and  $J = \langle x^2y, xy^4 \rangle$ . Compute the following ideals with SINGULAR:

- $I \cap J$ .
- $I \cdot J$ .
- $I : \langle x^3y^6 \rangle$ .
- $\sqrt{I}$ .
- Test if the polynomial  $x^7 + xy^8$  is in  $I$ .

Verify the results without SINGULAR.

**In-Class Exercise 3:** Welche der folgenden Ideale sind monomiale Ideale?

- $I = \langle x^2y - y^3, x^3 \rangle \triangleleft \mathbb{Q}[x, y, z]$ .
- $I = \langle x^4 - x^2y^2 + y^4, 2x^3 - xy^2, 2y^3 - x^2y \rangle \triangleleft \mathbb{Q}[x, y, z]$
- $I = \langle x12y7 + x9y + xyz3 + yz3, x8 - xyz, yz3, x8 - yz3, x12y7 \rangle \triangleleft \mathbb{Q}[x, y, z]$ .